arXiv:1301.5775v1 [math-ph] 24 Jan 2013

Comment on star-star relations in statistical mechanics and elliptic gamma-function identities

Vladimir V. Bazhanov[1,2], Andrew P. Kels[1] and Sergey M. Sergeev[3]

[1] *Department of Theoretical Physics, Research School of Physics and Engineering,*
Australian National University, Canberra, ACT 0200, Australia.

[2] *Mathematical Sciences Institute,*
Australian National University, Canberra, ACT 0200, Australia.

[3] *Faculty of Education Science Technology & Mathematics,*
University of Canberra, Bruce ACT 2601, Australia.

Abstract

We prove a recently conjectured star-star relation, which plays the role of an integrability condition for a class of 2D Ising-type models with multicomponent continuous spin variables. Namely, we reduce this relation to an identity for elliptic gamma functions, previously obtained by Rains.

1 Introduction

Recently two of us [1, 2] introduced a new class of exactly solvable 2D lattice models of statistical mechanics, which involve continuous spin variables taking values on a circle. The interest in these models is motivated by various applications. In statistical mechanics they serve as rather general "master models", containing many important particular limits, such as the Ising, chiral Potts [3, 4], Kashiwara-Miwa [5], Faddeev-Volkov [6, 7] and other models. Mathematically, the new models are deeply related to the theory of elliptic hypergeometric functions [8]. For instance, the celebrated elliptic beta integral [9], which lies at the basis of this theory, is shown [1] to be a Yang-Baxter (star-triangle) relation, defining perfectly physical integrable lattice models of statistical mechanics. Other interesting connections are discussed in [10–13]. We mention, in particular, that the elliptic gamma-functions arise in calculations of superconformal indices connected with electric-magnetic dualities in 4D $\mathcal{N} = 1$ superconformal Yang-Mills theories [14]. Most remarkably, as recently discovered in [15–17], the superconformal indices in 4D superconformal quiver gauge theories precisely coincide with partition functions of the 2D lattice "master models" [1, 2] discussed here. Interestingly, in this correspondence the Seiberg duality for the superconformal indices reduces to Baxter's Z-invariance [18] for the partition function under (generalized) "star-triangular moves" of the 2D lattice.

Here we resolve an outstanding question for these models concerning the so-called *star-star relation*, conjectured in [2]. This relation serves as an integrability condition, as it implies the Yang-Baxter equation, the Z-invariance of the partition function and the commutativity of row-to-row transfer matrices. In this letter we completely prove this star-star relation by reducing it to a transformation formula for elliptic hypergeometric integrals, previously obtained by Rains [19].

2 Edge-interaction model with continuous spins

In this Section we formulate the star-star relation conjectured in [2] (see (14) below). First, we need to briefly describe the associated two-dimensional edge-interaction model; full details can be found in [2]. Consider the regular square lattice, drawn diagonally as in Fig. 1. The edges of the lattice are shown with bold lines and the sites are shown with either open or filled circles in a checkerboard order. At the moment we will not distinguish these two type of sites. At each lattice site place a n-component continuous spin variable

$$\boldsymbol{x} = \{x_1, \ldots, x_n\} \in \mathbb{R}^n, \qquad 0 \leq x_j < \pi, \qquad \sum_{j=1}^{n} x_j = 0 \pmod{\pi} \ . \tag{1}$$

Note that due to the restriction on the total sum, there are only $(n-1)$ independent variables x_j. For further reference define the integration measure

$$\int d\boldsymbol{x} = \int_0^\pi \cdots \int_0^\pi dx_1 \cdots dx_{n-1} \ , \tag{2}$$

over the space of states of a single spin. Fig. 1 also shows an auxiliary *medial* lattice whose sites lie on the edges of the original square lattice. The medial lattice is drawn with alternating thin and dotted lines. The lines are directed as indicated by arrows. To each horizontal (vertical) line on the medial lattice assign a rapidity variable u (v). In general these variables may be different for different lines. However, a convenient level of generality that we shall use here is to assign the same rapidity u to all thin horizontal lines and the same variable u' to all dotted horizontal lines. Similarly, assign the variables v and v' to thin and dotted vertical lines as indicated in Fig. 1.



Figure 1: The square lattice shown with bold sites and bold edges drawn diagonally. The associated medial lattice is drawn with thin and dotted horizontal and vertical lines. The lines are oriented and carry rapidity variables u, u', v and v'.

Two spins interact only if they are connected by an edge. To define the Boltzmann weights we need to introduce the elliptic gamma-function [8]. Let q, p be two elliptic nomes (they play the role of the temperature-like parameters),

$$\mathsf{p} \;=\; \mathrm{e}^{\mathsf{i}\pi\sigma}\;, \quad \mathsf{q} = \mathrm{e}^{\mathsf{i}\pi\tau}\;, \quad \operatorname{Im}\sigma > 0, \quad \operatorname{Im}\tau > 0\,, \tag{3}$$

and

$$\eta = -\mathsf{i}\pi(\sigma+\tau)/2\,, \tag{4}$$

denote the "crossing parameter". Define the elliptic gamma-function[1]

$$\Phi(z) \;=\; \prod_{j,k=0}^{\infty} \frac{1-\mathrm{e}^{2\mathsf{i}z}\mathsf{q}^{2j+1}\mathsf{p}^{2k+1}}{1-\mathrm{e}^{-2\mathsf{i}z}\mathsf{q}^{2j+1}\mathsf{p}^{2k+1}} \;=\; \exp\left\{\sum_{k\neq 0}\frac{\mathrm{e}^{-2\mathsf{i}zk}}{k(\mathsf{q}^k-\mathsf{q}^{-k})(\mathsf{p}^k-\mathsf{p}^{-k})}\right\}\;, \tag{5}$$

where the product formula is valid for all z, while the exponential formula is only valid in the strip

$$-\operatorname{Re}\eta < \operatorname{Im} z < \operatorname{Re}\eta\;. \tag{6}$$

The function (5) possesses simple periodicity and "reflection" properties

$$\Phi(z+\pi) = \Phi(z)\,, \qquad \Phi(z)\,\Phi(-z) = 1\,. \tag{7}$$

Each edge is assigned a Boltzmann weight which depends on spins at the ends of the edge and on two rapidities passing through the edge. There are four types of edges differing by orientations and types of the directed rapidity lines passing through the edge. They are assigned different Boltzmann weights as shown in Fig.2. These weights are defined as

$$\mathbb{W}_\alpha(\boldsymbol{x},\boldsymbol{y}) = \kappa_n(\alpha)^{-1}\prod_{j,k=1}^{n}\Phi(x_j-y_k+\mathsf{i}\alpha), \quad \overline{\mathbb{W}}_\alpha(\boldsymbol{x},\boldsymbol{y}) = \sqrt{\mathbb{S}(\boldsymbol{x})\mathbb{S}(\boldsymbol{y})}\,\mathbb{W}_{\eta-\alpha}(x,y)\;, \tag{8}$$

---

[1] Our function $\Phi(z)$ coincides with $\Gamma(\mathrm{e}^{-2\mathsf{i}(z-\eta)};\mathsf{p}^2,\mathsf{q}^2)$ in the notation of ref. [8].


x
y
u
v'
$\mathbb{W}_{u-v'}(\boldsymbol{x},\boldsymbol{y})$
x
u
y
v
$\overline{\mathbb{W}}_{u-v}(\boldsymbol{y},\boldsymbol{x})$
y
x
u'
v
$\mathbb{W}_{u'-v}(\boldsymbol{x},\boldsymbol{y})$
y
u'
x
v'
$\overline{\mathbb{W}}_{u'-v'}(\boldsymbol{y},\boldsymbol{x})$

Figure 2: Four different types of edges and their Boltzmann weights.

where $\kappa_n(\alpha)$ is a normalization factor. The single-spin function $\mathbb{S}$ is given by

$$\mathbb{S}(\boldsymbol{x}) \;=\; \varkappa_s^{-1} \prod_{j\neq k} \Big\{ \Phi(x_j - x_k + \mathsf{i}\eta) \Big\}^{-1}, \qquad \varkappa_s \;=\; n! \left(\frac{\pi}{G(\mathsf{q})\, G(\mathsf{p})} \right)^{n-1}, \tag{9}$$

where the indices j, k run over the values $1, 2, \ldots, n$ and

$$G(z) = \prod_{k=1}^{\infty} (1 - z^{2k})\,. \tag{10}$$

The partition function is defined as

$$Z = \int \prod_{\langle \boldsymbol{x}\boldsymbol{y} \rangle} \mathbb{W}_{u-v'}(\boldsymbol{x},\boldsymbol{y}) \prod_{\langle \boldsymbol{y}\boldsymbol{x} \rangle} \overline{\mathbb{W}}_{u-v}(\boldsymbol{y},\boldsymbol{x}) \prod_{\langle \boldsymbol{x}\boldsymbol{y} \rangle} \mathbb{W}_{u'-v}(\boldsymbol{x}\boldsymbol{y}) \prod_{\langle \boldsymbol{y},\boldsymbol{x} \rangle} \overline{\mathbb{W}}_{u'-v'}(\boldsymbol{y},\boldsymbol{x}) \prod_{\text{sites}} d\boldsymbol{x}, \tag{11}$$

where the four products are taken, respectively, over the four types of edges shown in Fig. 2. The integral is taken over all configurations of the spin variables on the internal lattice sites. The boundary spins are kept fixed.

Note that the lattice in Fig. 1 can be formed by periodic translations of a *four-edge star*, consisting of four edges meeting at the same site. A little inspection shows that there are only two different types of such stars shown in Fig. 3. They are either centred around "white" sites, shown with open circles or around "black" sites, shown with filled circles. Applying the rules of Fig. 2 one



Figure 3: Two types of four-edge stars: a white-centred star $\mathbb{V}^{(1)}$ (left) and a black-centred star $\mathbb{V}^{(2)}$ (right).

can write Boltzmann weights corresponding to these stars

$$\mathbb{V}^{(1)}_{\boldsymbol{uv}}\left(\begin{array}{cc}\boldsymbol{a} & \boldsymbol{b}\\ \boldsymbol{c} & \boldsymbol{d}\end{array}\right)=\int d\boldsymbol{x}\,\overline{\mathbb{W}}_{u-v}(\boldsymbol{c},\boldsymbol{x})\,\overline{\mathbb{W}}_{u'-v'}(\boldsymbol{b},\boldsymbol{x})\,\mathbb{W}_{u'-v}(\boldsymbol{x},\boldsymbol{a})\,\mathbb{W}_{u-v'}(\boldsymbol{x},\boldsymbol{d})\;, \tag{12}$$

and

$$\mathbb{V}^{(2)}_{\boldsymbol{uv}}\left(\begin{array}{cc}\boldsymbol{a} & \boldsymbol{b}\\ \boldsymbol{c} & \boldsymbol{d}\end{array}\right)=\int d\boldsymbol{y}\,\overline{\mathbb{W}}_{u-v}(\boldsymbol{y},\boldsymbol{b})\,\overline{\mathbb{W}}_{u'-v'}(\boldsymbol{y},\boldsymbol{c})\,\mathbb{W}_{u'-v}(\boldsymbol{d},\boldsymbol{y})\,\mathbb{W}_{u-v'}(\boldsymbol{a},\boldsymbol{y})\;, \tag{13}$$

where the bold symbols $\boldsymbol{u}=[u,u']$ and $\boldsymbol{v}=[v,v']$ stand for the rapidity pairs. It turns out that the above two expressions are simply connected to each other

$$\mathbb{W}_{v'-v}(\boldsymbol{d},\boldsymbol{c})\,\mathbb{W}_{u'-u}(\boldsymbol{d},\boldsymbol{b})\;\mathbb{V}^{(1)}_{\boldsymbol{uv}}\left(\begin{array}{cc}\boldsymbol{a} & \boldsymbol{b}\\ \boldsymbol{c} & \boldsymbol{d}\end{array}\right)\;=\;\mathbb{W}_{v'-v}(\boldsymbol{b},\boldsymbol{a})\,\mathbb{W}_{u'-u}(\boldsymbol{c},\boldsymbol{a})\;\mathbb{V}^{(2)}_{\boldsymbol{uv}}\left(\begin{array}{cc}\boldsymbol{a} & \boldsymbol{b}\\ \boldsymbol{c} & \boldsymbol{d}\end{array}\right)\,. \tag{14}$$

This is precisely the *star-star relation* conjectured in [2] (in the same paper it was also verified in a few orders of perturbation theory in the parameters p and q). Below we will give a complete proof of (14) by reducing it to a mathematical identity, previously obtained by Rains [19]. Apparently, the above star-star relation is the simplest condition for the Boltzmann weights which ensures the integrability of the considered model[2]. In particular, it implies the commutativity of the row-to-row transfer matrices.

3 Proof of the star-star relation

Here we will use the standard notation for the elliptic gamma-function, which is simply related to our definition (5),

$$\Gamma(z;\mathsf{p}^2,\mathsf{q}^2)=\prod_{j,k=0}^{\infty}\frac{(1-\mathsf{p}^{2j+2}\,\mathsf{q}^{2k+2}/z)}{(1-\mathsf{p}^{2j}\,\mathsf{q}^{2k}\,z)}\,,\qquad \Phi(x)=\Gamma(\mathsf{pq}\,\mathrm{e}^{-2\mathrm{i}x};\mathsf{p}^2,\mathsf{q}^2)\,,\qquad \mathsf{pq}=\mathrm{e}^{-2\eta}\,. \tag{15}$$

In the following we will omit the nome arguments p^2 and q^2, assuming that $\Gamma(z)\equiv\Gamma(z;\mathsf{p}^2,\mathsf{q}^2)$. Following Rains [19] introduce the following elliptic hypergeometric integral[3]

$$\mathbf{I}^{(n-1)}_{A_{n-1}}(\{t_i\},\{s_i\})\;\stackrel{\mathrm{def}}{=}\;\frac{1}{n!}\left(\frac{G(\mathsf{p})\,G(\mathsf{q})}{2\pi\mathsf{i}}\right)^{n-1}\int_{|z_k|=1}\frac{\prod_{k=1}^{n}\prod_{j=1}^{2n}\Gamma(t_jz_k)\,\Gamma(s_j/z_k)}{\prod_{k\neq\ell}\Gamma(z_k/z_\ell)}\prod_{k=1}^{n-1}\frac{dz_k}{z_k}\,, \tag{16}$$

which involve $4n$ independent parameters

$$\{t_i\}=\{t_1,t_2,\ldots,t_{2n}\},\quad \{s_i\}=\{s_1,s_2,\ldots,s_{2n}\},\quad |t_i|,|s_i|<1,\quad i=1,\ldots,2n\,, \tag{17}$$

where $n\geq 2$ and the function $G(z)$ is defined in (10). The indices k and ℓ in the denominator of (16) run over the values $1,\ldots,n$. All integrations are taken over the unit circles $|z_k|=1$, and the variable z_n is determined by the constraint

$$z_1z_2\cdots z_n=1. \tag{18}$$

---

[2] For $n=2$ the star-star relation (14) is just a consequence of the star-triangle relation, Eq.(1.5) of [1], which is equivalent to the elliptic beta integral [9, 10]. However, for $n\geq 3$ the corresponding star-triangle relation apparently does not exist (at least it is not known to the authors) and the star-star relation (14) seems to be the simplest relation of this type.

[3] We follow Sect.4 of [19], where we set $Z\equiv 1$. Moreover, our indices numerating parameters in (17) start from 1 instead of 0 in [19].

The theorem 4.1 of [19] (where we set $m = n$ and $Z = 1$) states the following transformation formula,

$$\mathbf{I}^{(n-1)}_{A_{n-1}}(\{t_i\},\{s_i\}) = \Big(\prod_{j,k=1}^{2n}\Gamma(t_j s_k)\Big)\, \mathbf{I}^{(n-1)}_{A_{n-1}}(\{\widetilde{t}_i\},\{\widetilde{u}_i\}) \tag{19}$$

where the new parameters $\{\widetilde{t}_i\}$ and $\{\widetilde{s}_i\}$ in the RHS are given by

$$\widetilde{t}_i = T^{\frac{1}{n}}\, t_i^{\,-1}, \qquad \widetilde{u}_i = U^{\frac{1}{n}}\, s_i^{-1}, \qquad T = \prod_{j=1}^{2n} t_j, \qquad U = \prod_{j=1}^{2n} s_j\,, \quad i = 1, 2, \ldots, 2n\,. \tag{20}$$

Consider now the star weight (12) and make a change of variables $z_j = \mathrm{e}^{+2\mathrm{i}x_j}$ and

$$t_j = \mathrm{e}^{-2(u-v)-2\mathrm{i}c_j}, \quad t_{n+j} = \mathrm{e}^{-2(u'-v')-2\mathrm{i}b_j}, \quad s_j = \mathrm{e}^{2(u'-v-\eta)+2\mathrm{i}a_j}\,, \quad s_{n+j} = \mathrm{e}^{2(u-v'-\eta)+2\mathrm{i}d_j}\,, \tag{21}$$

where a_j, b_j, c_j, d_j are the components of the spin variables and $j = 1, \ldots, n$. Note that due to (1) the new variables z_j obey the constraint (18). Now taking into account that (9) can be written as

$$\mathbb{S}(\boldsymbol{x}) = \kappa_s^{-1}\Big(\prod_{j\neq k}\Gamma(z_j/z_k)\Big)^{-1}, \qquad z_j = \mathrm{e}^{2\mathrm{i}x_j}, \tag{22}$$

it is not difficult to check that

$$\mathbb{V}^{(1)}_{\boldsymbol{uv}}\begin{pmatrix} \boldsymbol{a} & \boldsymbol{b} \\ \boldsymbol{c} & \boldsymbol{d}\end{pmatrix} \;=\; \varrho\, \mathbf{I}^{(n-1)}_{A_{n-1}}(\{t_i\},\{s_i\})\,, \tag{23}$$

where

$$\varrho = \frac{\sqrt{\mathbb{S}(\boldsymbol{c})\mathbb{S}(\boldsymbol{b})}}{\kappa_n(\eta-u+v)\kappa_n(\eta-u'+v')\kappa_n(u'-v)\kappa_n(u-v')}\,, \tag{24}$$

Further, using (15) and the reflection property (7) for the elliptic gamma-function, one can re-write the ratio of the $\mathbb{W}$-factors entering (14) in the form

$$\mathbb{W}_{v'-v}(\boldsymbol{b},\boldsymbol{a})\,\mathbb{W}_{u'-u}(\boldsymbol{c},\boldsymbol{a})\Big[\mathbb{W}_{v'-v}(\boldsymbol{d},\boldsymbol{c})\,\mathbb{W}_{u'-u}(\boldsymbol{d},\boldsymbol{b})\Big]^{-1} = \prod_{j,k=1}^{2n}\Gamma(t_j s_k) \tag{25}$$

Next, substituting (21) into (20) one gets

$$\widetilde{t}_j = \mathrm{e}^{-2(u'-v')+2\mathrm{i}c_j}\,, \quad \widetilde{t}_{n+j} = \mathrm{e}^{-2(u-v)+2\mathrm{i}b_j}\,, \quad \widetilde{s}_j = \mathrm{e}^{2(u-v'-\eta)-2\mathrm{i}a_j}\,, \quad \widetilde{s}_{n+j} = \mathrm{e}^{2(u'-v-\eta)-2\mathrm{i}d_j}\,. \tag{26}$$

Consider now the star weight (13) and make a change of variables $z_j = \mathrm{e}^{-2\mathrm{i}x_j}$ (note the minus sign in the exponent). Using the variables (26) one obtains

$$\mathbb{V}^{(2)}_{\boldsymbol{uv}}\begin{pmatrix} \boldsymbol{a} & \boldsymbol{b} \\ \boldsymbol{c} & \boldsymbol{d}\end{pmatrix} = \varrho\, \mathbf{I}^{(n-1)}_{A_{n-1}}(\{\widetilde{t}_i\},\{\widetilde{s}_i\}) \tag{27}$$

where ϱ is the same as in (24). The relations (23), (25) and (27) immediately imply that the star-star relation (14) is equivalent to the Rains transformation formula (19) for the elliptic hypergeometric integrals, obtained in [19].

Acknowledgments

We are grateful to Hjalmar Rosengren who brought our attention to the the work of Rains [19]. The work was partially supported by the Australian Research Council.

References

[1] V. V. Bazhanov and S. M. Sergeev, "*A Master solution of the quantum Yang-Baxter equation and classical discrete integrable equations*," `arXiv:1006.0651 [math-ph]`.

[2] V. V. Bazhanov and S. M. Sergeev, "*Elliptic gamma-function and multi-spin solutions of the Yang-Baxter equation*," Nucl.Phys. **B856** (2012) 475–496, `arXiv:1106.5874 [math-ph]`.

[3] R. J. Baxter, J. H. H. Perk, and H. Au-Yang, "*New solutions of the star triangle relations for the chiral Potts model*," Phys. Lett. **A128** (1988) 138–142.

[4] H. Au-Yang, B. M. McCoy, J. H. H. perk, S. Tang, and M.-L. Yan, "*Commuting transfer matrices in the chiral Potts models: Solutions of star-triangle equations with genus* > 1," Phys. Lett. **A123** (1987) 219–223.

[5] M. Kashiwara and T. Miwa, "*A class of elliptic solutions to the star-triangle relation*," Nuclear Physics B **275** (1986) 121–134.

[6] A. Y. Volkov and L. D. Faddeev, "*Yang-Baxterization of the quantum dilogarithm*," Zapiski Nauchnykh Seminarov POMI **224** (1995) 146–154. English translation: J. Math. Sci. **88** (1998) 202-207.

[7] V. V. Bazhanov, V. V. Mangazeev, and S. M. Sergeev, "*Faddeev-Volkov solution of the Yang-Baxter Equation and Discrete Conformal Symmetry*," Nucl. Phys. **B784** (2007) 234–258, `hep-th/0703041`.

[8] V. P. Spiridonov, "*Essays on the theory of elliptic hypergeometric functions*," Uspekhi Mat. Nauk **63** (2008) no. 3(381), 3–72. `http://dx.doi.org/10.1070/RM2008v063n03ABEH004533`.

[9] V. P. Spiridonov, "*On the elliptic beta function*," Uspekhi Mat. Nauk **56 (1)** (2001) no. 1, 181–182. `http://dx.doi.org/10.1070/rm2001v056n01ABEH000374`. English translation: Russ. Math. Surveys **56** (1) (2001) 185-186.

[10] V. P. Spiridonov, "*Elliptic beta integrals and solvable models of statistical mechanics*," `arXiv:1011.3798 [hep-th]`.

[11] R. Kashaev, F. Luo, and G. Vartanov, "*A TQFT of Turaev-Viro type on shaped triangulations*," `arXiv:1210.8393 [math.QA]`.

[12] D. Chicherin, S. Derkachov, and A. Isaev, "*Conformal group: R-matrix and star-triangle relation*," `arXiv:1206.4150 [math-ph]`.

[13] D. Chicherin, S. Derkachov, D. Karakhanyan, and R. Kirschner, "*Baxter operators with deformed symmetry*," Nucl.Phys. **B868** (2013) 652–683, `arXiv:1211.2965 [math-ph]`.

[14] V. Spiridonov and G. Vartanov, "*Elliptic hypergeometry of supersymmetric dualities II. Orthogonal groups, knots, and vortices*," `arXiv:1107.5788 [hep-th]`.

[15] M. Yamazaki, "*Quivers, YBE and 3-manifolds*," JHEP **1205** (2012) 147, `arXiv:1203.5784 [hep-th]`.

[16] Y. Terashima and M. Yamazaki, "*Emergent 3-manifolds from 4d Superconformal Indices*," Phys.Rev.Lett. **109** (2012) 091602, `arXiv:1203.5792 [hep-th]`.

[17] D. Xie and M. Yamazaki, "*Network and Seiberg Duality*," JHEP **1209** (2012) 036, `arXiv:1207.0811 [hep-th]`.

[18] R. J. Baxter, "*Solvable eight-vertex model on an arbitrary planar lattice*," Philos. Trans. Roy. Soc. London Ser. A **289** (1978) no. 1359, 315–346.

[19] E. M. Rains, "*Transformations of elliptic hypergeometric integrals*," Ann. of Math. (2) **171** (2010) no. 1, 169–243. `http://dx.doi.org.virtual.anu.edu.au/10.4007/annals.2010.171.169`.